

# On a two-sided Turán problem

Dhruv Mubayi \*    Yi Zhao †

Department of Mathematics, Statistics, and Computer Science  
University of Illinois at Chicago  
Chicago, IL 60607

August 21, 2003

## Abstract

Given positive integers  $n, k, t$ , with  $2 \leq k \leq n$ , and  $t < 2^k$ , let  $m(n, k, t)$  be the minimum size of a family  $\mathcal{F}$  of nonempty subsets of  $[n]$  such that every  $k$ -set in  $[n]$  contains at least  $t$  sets from  $\mathcal{F}$ , and every  $(k-1)$ -set in  $[n]$  contains at most  $t-1$  sets from  $\mathcal{F}$ . Sloan et. al [11] determined  $m(n, 3, 2)$  and Füredi et. al [6] studied  $m(n, 4, t)$  for  $t = 2, 3$ . We consider  $m(n, 3, t)$  and  $m(n, 4, t)$  for all the remaining values of  $t$  and obtain their exact values except for  $k = 4$  and  $t = 6, 7, 11, 12$ . For example, we prove that  $m(n, 4, 5) = \binom{n}{2} - 17$  for  $n \geq 160$ . The values of  $m(n, 4, t)$  for  $t = 7, 11, 12$  are determined in terms of well-known (and open) Turán problems for graphs and hypergraphs. We also obtain bounds of  $m(n, 4, 6)$  that differ by absolute constants.

Keywords: Extremal problem, Hypergraph, Turán number, set system

MR Subject Classifications: 05D05, 05C35, 05C65

## 1 Introduction

We consider an extremal problem for set systems. Given integers  $n, k, t$ , with  $2 \leq k \leq n$ , and  $t < 2^k$ , a family  $\mathcal{F} \subset 2^{[n]} \setminus \emptyset$  is a  $(k, t)$ -system of  $[n]$  if every  $k$ -set in  $[n]$  contains at least  $t$  sets

---

\*Research supported in part by NSF grant DMS-9970325.

†Research supported in part by NSF grant DMS-9983703, a VIGRE Postdoctoral Fellowship at University of Illinois at Chicago.

from  $\mathcal{F}$ , and every  $(k-1)$ -set in  $[n]$  contains at most  $t-1$  sets from  $\mathcal{F}$ . Let  $m(n, k, t)$  denote the minimum size of a  $(k, t)$ -system of  $[n]$ . This threshold function first arose in problems on computer science [10, 11] (although the notation  $m(n, k, t)$  was not used until [6]). It was shown in [11] that  $m(n, k, t) = \Theta(n^{k-1})$  for  $1 < t < k$  and  $m(n, 3, 2) = \binom{n-1}{2} + 1$ . In [6],  $m(n, 4, 3)$  was determined exactly for large  $n$  and it was shown that for fixed  $k$ ,  $m(n, k, 2) = (1 + o(1))T_{k-1}(n, k, 2)$ , where  $T_r(n, k, t)$  is the generalized Turán number. For fixed  $k$  and  $t < 2^k$ , the order of magnitude of  $m(n, k, t)$  was determined in [9]. A special case of this result is the following proposition, where  $\binom{a}{\leq b} = \sum_{i=1}^b \binom{a}{i}$ .

**Proposition 1.** [9]  $m(n, k, 1) = \binom{n}{k}$ ,  $m(n, k, k) = n$ ,  $m(n, k, 2^k - 2) = \binom{n}{\leq k-1}$  and  $m(n, k, 2^k - 1) = \binom{n}{\leq k}$ .

In this paper we study  $m(n, k, t)$  for  $k = 3, 4$ . The case  $k = 3$  is not very difficult: Proposition 1 determines  $m(n, 3, t)$  for  $t \in \{1, 3, 6, 7\}$  and [11] shows that  $m(n, 3, 2) = \binom{n-1}{2} + 1$ . The remaining cases  $t = 4$  and  $t = 5$  are covered below.

**Proposition 2.**

$$m(n, 3, t) = \begin{cases} n + \binom{n}{2} - \lfloor n^2/4 \rfloor & t = 4, \\ n + \binom{n}{2} - \lfloor n/2 \rfloor & t = 5. \end{cases}$$

The main part of this paper is devoted to  $m(n, 4, t)$ , a problem which is substantially more difficult than the case  $k = 3$ . As mentioned above, both  $m(n, 4, 2)$  and  $m(n, 4, 3)$  were studied in [6]. It was shown in [11] how these two functions apply to frequent sets of Boolean matrices, a concept used in knowledge discovery and data mining. Perhaps the determination of  $m(n, 4, t)$  for other  $t$  will have similar applications.

The cases  $t = 1, 4, 14, 15$  are answered by Proposition 1 immediately. In this paper we obtain the exact values of  $m(n, 4, t)$  for  $t = 5, 8, 9, 10, 13$ . Our bounds for  $m(n, 4, 6)$  differ only by an absolute constant. For  $t = 7, 11, 12$ , we determine  $m(n, 4, t)$  exactly in terms of well-known (and open) Turán problems in extremal graph and hypergraph theory. Perhaps this connection provides additional motivation for investigating  $m(n, k, t)$  (the first connection between  $m(n, k, t)$  and Turán numbers was shown in [6] via  $m(n, k, 2) = (1 + o(1))T_{k-1}(n, k, 2)$ ).

For a family of  $r$ -uniform hypergraphs  $\mathcal{H}$ , let  $\text{ex}(n, \mathcal{H})$  be the maximum number of edges in an  $n$  vertex  $r$ -uniform hypergraph  $\mathcal{G}$  containing no member of  $\mathcal{H}$  as a subhypergraph. The (2-uniform) cycle of length  $l$  is written  $C_l$ . The complete 3-uniform hypergraph on four points is  $K_4^{(3)}$ , and the 3-uniform hypergraph on four points with three edges is  $H(4, 3)$ .

An  $(n, 3, 2)$ -packing is a 3-uniform hypergraph on  $n$  vertices such that every pair of vertices is contained in at most one edge. The packing number  $P(n, 3, 2)$  is the size of a maximal  $(n, 3, 2)$ -packing. Note that the maximal packing is a Steiner system when  $n \equiv 1$  or  $3 \pmod{6}$ .

**Theorem 3 (Main Theorem).**

$$m(n, 4, 5) = \binom{n}{2} - 17,$$

when  $n \geq 160$  and

$$\binom{n}{2} - 190 < m(n, 4, 6) \leq \binom{n}{2} - 5,$$

when  $n \geq 8$ . Furthermore,

$$\begin{aligned} m(n, 4, 7) &= n + \binom{n}{2} - \text{ex}(n, \{C_3, C_4\}), \\ m(n, 4, 8) &= n + \binom{n}{2} - 2n/3, \\ m(n, 4, 9) &= n + \binom{n}{2} - 1, \\ m(n, 4, 10) &= n + \binom{n}{2}, \\ m(n, 4, 11) &= n + \binom{n}{2} + \binom{n}{3} - \text{ex}(n, K_4^{(3)}), \\ m(n, 4, 12) &= n + \binom{n}{2} + \binom{n}{3} - \text{ex}(n, H(4, 3)), \\ m(n, 4, 13) &= n + \binom{n}{2} + \binom{n}{3} - P(n, 3, 2). \end{aligned}$$

It is worth recalling the known results for the three Turán numbers and the packing number  $P(n, 3, 2)$  in Theorem 3 above.

- It is known that  $(\frac{1}{2\sqrt{2}} + o(1))n^{3/2} \leq \text{ex}(n, \{C_3, C_4\}) \leq (\frac{1}{2} + o(1))n^{3/2}$  (Erdős-Rényi [3], Kővari-Sós-Turán [7]). Erdős and Simonovits [4] conjectured that  $\text{ex}(n, \{C_3, C_4\}) = (\frac{1}{2\sqrt{2}} + o(1))n^{3/2}$ .
- It is known that  $(5/9)\binom{n}{3} \leq \text{ex}(n, K_4^{(3)}) \leq (0.592 + o(1))\binom{n}{3}$  (Turán [14], Chung-Lu [2]). It was conjectured [14] that the lower bound is correct (Erdős offered \$1000 for a proof).
- It is known  $(2/7 + o(1))\binom{n}{3} \leq \text{ex}(n, H(4, 3)) \leq (1/3 - 10^{-6} + o(1))\binom{n}{3}$  (Frankl-Füredi [5], Mubayi [8]). It was conjectured [8] that  $\text{ex}(n, H(4, 3)) = (2/7 + o(1))\binom{n}{3}$ .
- Spencer [12] determine  $P(n, 3, 2)$  exactly:

$$P(n, 3, 2) = \begin{cases} \lfloor \frac{n}{3} \lfloor \frac{n-1}{2} \rfloor \rfloor - 1 & \text{if } n \equiv 5 \pmod{6}, \\ \lfloor \frac{n}{3} \lfloor \frac{n-1}{2} \rfloor \rfloor & \text{otherwise.} \end{cases}$$

This paper is organized as follows. In Section 2 we describe the main idea in the proofs and prove Propositions 2. The Main Theorem (Theorem 3) is proved in Section 3.

Most of our notations are standard:  $[n] = \{1, 2, \dots, n\}$ . For a set system  $\mathcal{F}$ , let  $\mathcal{F}_t$  denote the family of  $t$ -sets in  $\mathcal{F}$ , let  $\mathcal{F}_{\leq t} = \cup_{i \leq t} \mathcal{F}_i$  and  $\mathcal{F}_{\geq t} = \cup_{i \geq t} \mathcal{F}_i$ . If  $a \in \mathcal{F}$  and  $b \notin \mathcal{F}$ , we simply write  $\mathcal{F} - a$  for  $\mathcal{F} \setminus \{a\}$  and  $\mathcal{F} + b$  for  $\mathcal{F} \cup \{b\}$ . Given a set  $X$  and an integer  $a$ , let  $2^X = \{S : S \subseteq X\}$ ,  $\binom{X}{a} = \{S \subset X : |S| = a\}$ ,  $\binom{X}{\leq a} = \{S \subset X : 1 \leq |S| \leq a\}$  and  $\binom{X}{\geq a} = \{S \subset X : |S| \geq a\}$ . We write  $\mathcal{F}(X)$  for  $\mathcal{F} \cap 2^X$ . An  $r$ -graph on  $X$  is a (hyper)graph whose edges are  $r$ -subsets of  $X$ . All sets or subsets considered in this paper are nonempty unless specified differently.

## 2 Ideas in the proofs and $m(n, 3, t)$

In this section we make some basic observations on  $m(n, k, t)$  and prove Propositions 2.

Recall that a  $(k, t)$ -system  $\mathcal{F} \subseteq 2^{[n]} \setminus \emptyset$  satisfies the following two conditions:

**Property D (DENSE):** Every  $k$ -set in  $[n]$  contains at least  $t$  sets from  $\mathcal{F}$ ,

**Property S (SPARSE):** Every  $(k - 1)$ -set in  $[n]$  contains at most  $t - 1$  sets from  $\mathcal{F}$ .

The main idea in our proofs is to work with optimal  $(k, t)$ -systems which are defined as follows.

**Definition 4.** *Suppose that  $\mathcal{F}$  is a  $(k, t)$ -system of  $[n]$ . We say that  $\mathcal{F}$  is optimal if  $|\mathcal{F}| = m(n, k, t)$  and  $\sum_{S \in \mathcal{F}} |S|$  is minimal among all  $(k, t)$ -system of  $[n]$  with size  $m(n, k, t)$ .*

The advantage of considering optimal  $(k, t)$ -systems  $\mathcal{F}$  is that it allows us to assume certain structure on  $\mathcal{F}$ : if  $\mathcal{F}$  does not have such a structure, we always modify  $\mathcal{F}$  to  $\mathcal{F}'$  such that  $\mathcal{F}'$  is a  $(k, t)$ -system with  $\sum_{S \in \mathcal{F}'} |S| < \sum_{S \in \mathcal{F}} |S|$ , a contradiction to the optimality of  $\mathcal{F}$ . A typical modification of  $\mathcal{F}$  is replacing a set in  $\mathcal{F}$  by one of its subsets. Because the new system still satisfies Property **D**, we only need to check Property **S** in this case.

For example, if  $\mathcal{F}$  is an optimal  $(k, t)$ -system for  $t \geq 2^{k-1}$ , then we may assume that

$$A \in \mathcal{F} \Rightarrow (2^A \setminus \emptyset) \subset \mathcal{F}. \quad (\star)$$

Indeed, if  $A \in \mathcal{F}$  has a nonempty subset  $B \notin \mathcal{F}$ , then  $\mathcal{F}' = \mathcal{F} - A + B$  is also a  $(k, t)$ -system, because Property **S** holds trivially (any  $(k - 1)$ -set of  $[n]$  has at most  $2^{k-1} - 1 \leq t - 1$

nonempty subsets). Since  $\sum_{S \in \mathcal{F}'} |S| < \sum_{S \in \mathcal{F}} |S|$ , this contradicts the optimality of  $\mathcal{F}$ .

Now we consider  $m(n, 3, t)$  for  $3 \leq t \leq 7$ . Applying Proposition 1 directly, we have  $m(n, 3, 3) = n$ ,  $m(n, 3, 6) = \binom{n}{\leq 2}$  and  $m(n, 3, 7) = \binom{n}{\leq 3}$ .

**Proof of Proposition 2.** We determine  $m(n, 3, t)$  exactly for  $t = 4, 5$ . Recall that  $\mathcal{F}(S) = \mathcal{F} \cap 2^S$  for a set system  $\mathcal{F}$  and a set  $S$ .

Let  $\mathcal{F}$  be an optimal  $(3, t)$ -system with  $4 \leq t \leq 5$ . Since  $t \geq 4 \geq 2^2$ , we may assume that  $(\star)$  holds in  $\mathcal{F}$ . First, we claim that  $\binom{[n]}{1} \subset \mathcal{F}$ . Suppose instead, that there exists some  $a \in [n]$  such that  $\{a\} \notin \mathcal{F}$ . Pick a 3-set  $T = \{a, b, c\}$ . Since  $\{a\} \notin \mathcal{F}$ , by  $(\star)$ , we know that  $\mathcal{F}$  does not contain  $\{a, b\}, \{a, c\}$  and  $T$  as well. Thus  $|\mathcal{F}(T)| \leq 3$ , a contradiction to Property **D**. Second, we claim that  $\mathcal{F} \subset \binom{[n]}{\leq 2}$ . Suppose instead, that there exists a set  $T \in \mathcal{F}_3$ . Then  $|\mathcal{F}(T)| = 7$  by  $(\star)$  and consequently  $\mathcal{F}' = \mathcal{F} - T$  is a  $(3, t)$ -system of cardinality  $|\mathcal{F}| - 1$ , contradicting the optimality of  $\mathcal{F}$ .

When  $t = 4$ ,  $\mathcal{F}_2 = \mathcal{F} \setminus \binom{[n]}{1}$  is the edge set of a graph on  $n$  vertices in which every set of 3 vertices has at least one edge, *i.e.*,  $\overline{\mathcal{F}_2}$  is a  $K_3$ -free graph. Thus  $|\mathcal{F}_2| \geq \binom{n}{2} - \text{ex}(n, K_3) = \binom{n}{2} - \lfloor n^2/4 \rfloor$ . Consequently  $m(n, 3, 4) = n + |\mathcal{F}_2| \geq n + \binom{n}{2} - \lfloor n^2/4 \rfloor$ . On the other hand,  $\binom{[n]}{1} \cup E(G)$  is a  $(3, 4)$ -system, where  $G$  is a complete bipartite graph with two color classes of size  $\lfloor n/2 \rfloor$  and  $\lceil n/2 \rceil$ . Consequently  $m(n, 3, 4) = n + \binom{n}{2} - \lfloor n^2/4 \rfloor$ .

When  $t = 5$ ,  $\mathcal{F}_2 = \mathcal{F} \setminus \binom{[n]}{1}$  is the edge set of a graph on  $n$  vertices in which every 3 vertices have at least two edges. Therefore  $\overline{\mathcal{F}_2}$  is a matching  $M$  and  $|\mathcal{F}_2| = \binom{n}{2} - |M| \geq \binom{n}{2} - \lfloor n/2 \rfloor$ . Consequently  $m(n, 3, 5) \geq n + \binom{n}{2} - \lfloor n/2 \rfloor$  and equality holds for the  $(3, 5)$ -system  $\mathcal{F} = \binom{[n]}{1} \cup E(G)$ , where  $G$  is a complete graph except for a matching of size  $\lfloor n/2 \rfloor$ .  $\square$

### 3 The values of $m(n, 4, t)$

Applying Proposition 1, we obtain that  $m(n, 4, 1) = \binom{n}{4}$ ,  $m(n, 4, 4) = n$ ,  $m(n, 4, 14) = \binom{n}{\leq 3}$  and  $m(n, 4, 15) = \binom{n}{\leq 4}$ . In this section we prove Theorem 3, *i.e.*, determine  $m(n, 4, t)$  for  $5 \leq t \leq 13$ . We consider the cases  $7 \leq t \leq 13$  in Section 3.1. The more difficult cases  $t = 5, 6$  are studied in Section 3.2 and 3.3, respectively.

### 3.1 The cases $7 \leq t \leq 13$

Our proof is facilitated by the following four lemmas, whose proofs are postponed to the end of this section.

In Lemmas 5 - 8,  $\mathcal{F}$  is an optimal  $(4, t)$ -system.

**Lemma 5.** *If  $2 \leq t \leq 14$ , then  $\mathcal{F}_4 = \emptyset$ .*

**Lemma 6.** *If  $7 \leq t \leq 10$ , then  $\mathcal{F}_3 = \emptyset$ .*

**Lemma 7.** *If  $7 \leq t \leq 14$ , then  $\binom{[n]}{1} \subset \mathcal{F}$ .*

**Lemma 8.** *If  $11 \leq t \leq 14$ , then  $\binom{[n]}{2} \subset \mathcal{F}$ .*

**Proof of Theorem 3 for  $7 \leq t \leq 13$ :**

By Lemmas 5, 6 and 7, we conclude that

$$\binom{[n]}{1} \subset \mathcal{F} \subset \binom{[n]}{\leq 2} \quad \text{for } 7 \leq t \leq 10.$$

Clearly, when  $t = 10$ ,  $\mathcal{F} = \binom{[n]}{\leq 2}$  and consequently  $m(n, 4, 10) = \binom{n}{\leq 2}$ .

When  $t = 9$ ,  $\mathcal{F}_2 = \mathcal{F} \setminus \binom{[n]}{1}$  is the edge set of a graph on  $[n]$  in which every 4-set has at least 5 edges. Then there is at most one edge absent from  $\mathcal{F}_2$ , or  $|\mathcal{F}_2| \geq \binom{n}{2} - 1$ . Consequently  $m(n, 4, 9) \geq n + \binom{n}{2} - 1$  and equality holds when  $\mathcal{F} = \binom{[n]}{\leq 2} \setminus e$  for some  $e \in \binom{[n]}{2}$ .

When  $t = 8$ ,  $\mathcal{F}_2 = \mathcal{F} \setminus \binom{[n]}{1}$  is the edge set of a graph on  $[n]$  in which every 4-set has at least 4 edges. Therefore,  $\overline{\mathcal{F}_2}$  contains no  $K_3$ ,  $S_3$  (a star with 3 leaves), or  $P_3$  (a path of length 3). Thus all connected components of  $\overline{\mathcal{F}_2}$  have size at most 3 and each component is either an edge or  $P_2$ . So  $|\overline{\mathcal{F}_2}| \leq \lfloor 2n/3 \rfloor$  and  $|\mathcal{F}| \geq n + \binom{n}{2} - \lfloor 2n/3 \rfloor$ . Consequently  $m(n, 4, 8) = n + \binom{n}{2} - \lfloor 2n/3 \rfloor$  and the optimal system is  $\binom{[n]}{\leq 2} \setminus E(G)$ , where  $G$  is the union of disjoint copies of  $P_2$  and  $P_1$  covering  $[n]$  with maximum copies of  $P_2$ .

When  $t = 7$ ,  $\mathcal{F}_2 = \mathcal{F} \setminus \binom{[n]}{1}$  is the edge set of a graph on  $[n]$  in which every 4-set has at least 3 edges. Let  $G$  be a graph on  $[n]$  with  $E(G) = \overline{\mathcal{F}_2}$ . Then  $G$  contains no copies of  $C_4$  or  $C_3^+$  ( $C_3$  plus an edge). If  $C_3$  is also absent in  $G$ , then  $e(G) \leq ex(n, \{C_3, C_4\})$ . Otherwise, assume that  $G$  contains  $t (\geq 1)$  copies of  $C_3$  on a vertex-set  $T$ . Because  $G$  is  $C_3^+$ -free, the copies of  $C_3$  must be vertex-disjoint and

$$e(G) = 3t + e(G \setminus T) \leq 3t + ex(n - 3t, \{C_3, C_4\}) \leq ex(n, \{C_3, C_4\}),$$

where the last inequality is an easy exercise. Consequently  $m(n, 4, 7) \geq n + \binom{n}{2} - ex(n, \{C_3, C_4\})$  and equality holds when  $\mathcal{F} = \binom{[n]}{\leq 2} \setminus E(G)$ , where  $G$  is an extremal graph without  $C_3$  or  $C_4$ .

By Lemma 5, 7 and 8, we conclude that

$$\binom{[n]}{\leq 2} \subset \mathcal{F} \subset \binom{[n]}{\leq 3} \quad \text{for } 11 \leq t \leq 13.$$

When  $t = 11$ ,  $\mathcal{F}_3 = \mathcal{F} \setminus \binom{[n]}{\leq 2}$  is the edge set of a 3-graph in which every 4-set has at least one hyper-edge. In other words, the 3-graph  $([n], \overline{\mathcal{F}}_3)$  contains no  $K_4^{(3)}$  and therefore  $|\overline{\mathcal{F}}_3| \leq ex(n, K_4^{(3)})$ . Consequently  $m(n, 4, 11) \geq \binom{n}{\leq 3} - ex(n, K_4^{(3)})$  and equality holds when  $\mathcal{F} = \binom{[n]}{\leq 3} \setminus \mathcal{H}$ , where  $\mathcal{H}$  is the edge set of an extremal 3-graph without  $K_4^{(3)}$ .

By a similar argument, we obtain that  $m(n, 4, 12) \geq \binom{n}{\leq 3} - ex(n, H(4, 3))$  and equality holds when  $\mathcal{F} = \binom{[n]}{\leq 3} \setminus \mathcal{H}$ , where  $\mathcal{H}$  is the edge set of an extremal 3-graph without  $H(4, 3)$ .

Finally, when  $t = 13$ ,  $\overline{\mathcal{F}}_3$  is an  $(n, 3, 2)$ -packing since every 4-set of  $[n]$  contains at most one hyper-edge of  $\overline{\mathcal{F}}_3$ . Since  $|\overline{\mathcal{F}}_3| \leq P(n, 3, 2)$ , we have  $m(n, 4, 13) \geq \binom{n}{\leq 3} - P(n, 3, 2)$  and equality holds when  $\overline{\mathcal{F}}_3$  is a maximal  $(n, 3, 2)$ -packing.  $\square$

Before verifying Lemma 5, we start with a technical lemma, which is very useful in the cases  $5 \leq t \leq 7$ .

**Lemma 9.** *Suppose that  $t \in \{5, 6, 7\}$  and  $\mathcal{F}$  is an optimal  $(4, t)$ -system. Fix a set  $P \in \binom{[n]}{\leq 2} \setminus \mathcal{F}$  and let*

$$\mathcal{T} = \{T \in \binom{[n]}{3} : T \supset P, |\mathcal{F}(T)| = t - 1\}. \quad (1)$$

*If  $\mathcal{T} \subset \mathcal{F}$ , then  $T \notin \mathcal{F}$  for every 3-set  $T \supset P$ .*

**Proof.** Suppose instead, that there exists a 3-set  $T_0 \supset P$  and  $T_0 \in \mathcal{F}$ . If  $\mathcal{T} = \emptyset$ , then let  $\mathcal{F}' = \mathcal{F} - T_0 + P$ . It is clear that  $\mathcal{F}'$  satisfies Property **D**.  $\mathcal{F}'$  also satisfies Property **S** because  $|\mathcal{F}'(Y)| = |\mathcal{F}(Y)| + 1 \leq t - 2 + 1 = t - 1$  for every 3-set  $Y \supset P$ . Therefore  $\mathcal{F}'$  is a  $(4, t)$ -system, a contradiction to the optimality of  $\mathcal{F}$ .

Now assume that  $\mathcal{T} \neq \emptyset$ . We claim that  $\mathcal{F}' = \mathcal{F} - \mathcal{T} + P$  is a  $(4, t)$ -system, contradicting the optimality of  $\mathcal{F}$ . To check Property **D**, we only need to consider those 4-sets  $S$  which contain two members  $T_1, T_2$  of  $\mathcal{T}$  (because  $|\mathcal{F}'(Q)| = |\mathcal{F}(Q)|$  for every 4-set  $Q$  that contains at most one member of  $\mathcal{T}$ ). Since  $|\mathcal{F}'(S)| \geq |\mathcal{F}(T_1)| + |\mathcal{F}(T_2)| - |\mathcal{F}(P)| \geq 2(t - 1) - 2 = 2t - 4 \geq t + 1$

(using the assumption that  $t \geq 5$ ), we have  $|\mathcal{F}'(S)| \geq t + 1 - 2 + 1 = t$ . On the other hand,  $\mathcal{F}'$  also satisfies Property **S** since for every 3-set  $Y \supset P$ ,  $|\mathcal{F}'(Y)| = |\mathcal{F}(Y)| = t - 1$  if  $Y \in \mathcal{T}(\subset \mathcal{F})$ , otherwise  $|\mathcal{F}'(Y)| = |\mathcal{F}(Y)| + 1 \leq t - 2 + 1 = t - 1$ .  $\square$

**Proof of Lemma 5.** We are to show that  $\mathcal{F}_4 = \emptyset$  for  $2 \leq t \leq 14$ .

When  $8 \leq t \leq 14$ ,  $(\star)$  holds in  $\mathcal{F}$  (since  $t \geq 2^3$ ). We may thus assume that  $\mathcal{F}$  contains no 4-set, otherwise removing these 4-sets results in a smaller  $(4, t)$ -system, a contradiction to the optimality of  $\mathcal{F}$ .

Let  $2 \leq t \leq 7$ . Suppose to the contrary, that there exists a set  $S \in \mathcal{F}_4$ . We may assume that  $|\mathcal{F}(S)| = t$ , otherwise  $S$  could be removed from  $\mathcal{F}$ . Let  $\mathcal{T} = \binom{S}{3} \setminus \mathcal{F}$ .

**Case 1.**  $\mathcal{T} \neq \emptyset$ .

Suppose that  $T_0 \in \mathcal{T}$  has the minimal value of  $|\mathcal{F}(T)|$  among all  $T \in \mathcal{T}$ . We claim that  $|\mathcal{F}(T_0)| \leq t - 2$ . Suppose instead, that  $|\mathcal{F}(T_0)| \geq t - 1$ . If  $|\mathcal{T}| < 4$ , then there exists  $T_1 \in \binom{S}{3} \cap \mathcal{F}$ . Because  $T_1, S \in \mathcal{F}$ , we have  $|\mathcal{F}(S)| \geq |\mathcal{F}(T_0)| + 2 \geq t - 1 + 2 > t$ , a contradiction to the assumption that  $|\mathcal{F}(S)| = t$ . If  $|\mathcal{T}| = 4$ , then for every  $T \in \binom{S}{3}$ , we have  $|\mathcal{F}(T)| \geq t - 1$  and  $T \notin \mathcal{F}$ . Since  $|\cup_{T \in \binom{S}{3}} \mathcal{F}(T)| = |\mathcal{F}(S) \setminus S| = t - 1$ , we have  $\mathcal{F}(T_1) = \mathcal{F}(T_2) \neq \emptyset$  for every  $T_1, T_2 \in \binom{S}{3}$ . But this is impossible because  $\cap_{i=1}^4 T_i = \emptyset$ . Now let  $\mathcal{F}' = \mathcal{F} - S + T_0$ . Trivially  $\mathcal{F}'$  satisfies Property **D** and because  $|\mathcal{F}'(T_0)| \leq t - 1$ ,  $\mathcal{F}'$  satisfies Property **S** as well. Thus  $\mathcal{F}'$  is a  $(4, t)$ -system, a contradiction to the optimality of  $\mathcal{F}$ .

**Case 2.**  $\mathcal{T} = \emptyset$ , i.e.,  $\binom{S}{3} \subset \mathcal{F}$ .

Note that this case does not exist for  $t = 2, 3, 4$ , because it implies that  $|\mathcal{F}(S)| \geq 4 + 1$ , a contradiction to the assumption  $|\mathcal{F}(S)| = t$ .

When  $t = 5$ , we know that  $\mathcal{F}(S) = \{S\} \cup \binom{S}{3}$ . Pick any two elements  $a, b \in S$  and consider  $\mathcal{T} = \{\{a, b, c\} : |\mathcal{F}(\{a, b, c\})| = 4\}$ . Since  $\mathcal{F}(\{a, b\}) = \emptyset$ , it must be the case that  $\mathcal{F}(T) = \{\{c\}, \{c, a\}, \{c, b\}, \{c, a, b\}\}$  for every  $T = \{a, b, c\} \in \mathcal{T}$ . In particular,  $\mathcal{T} \subset \mathcal{F}$ . We may therefore apply Lemma 9 to conclude that  $T \notin \mathcal{F}$  for every 3-set  $T \subset \{a, b\}$ . This is a contradiction to the assumption that  $T \in \mathcal{F}$  for all  $T \in \binom{S}{3}$ .

When  $t = 6, 7$ , since  $|\mathcal{F}(S)| \leq 7$  and  $\binom{S}{\geq 3} \subset \mathcal{F}$ , we have  $|\mathcal{F} \cap \binom{S}{\leq 2}| \leq 2$ . Consequently there exist  $a, b \in S$  such that  $\mathcal{F}(\{a, b\}) = \emptyset$ . Since  $\mathcal{T} = \{\{a, b, c\} : |\mathcal{F}(\{a, b, c\})| = t\} = \emptyset$ , we may again apply Lemma 9 and derive a contradiction as in the previous paragraph.  $\square$

**Proof of Lemma 6.** We are to show that  $\mathcal{F}_3 = \emptyset$  for  $7 \leq t \leq 10$ . Suppose to the contrary, that there exists a set  $T \in \mathcal{F}_3$ . We now separate the case  $t = 7$  and the cases  $t = 8, 9, 10$ .

**Case 1.**  $t = 7$ .

Since  $|\mathcal{F}(T)| < 7$  (by Property **S**), there exists a set  $P \in \binom{T}{\leq 2} \setminus \mathcal{F}$ . Define  $\mathcal{T}$  as in (1), trivially  $\mathcal{T} \subset \mathcal{F}$ . We may apply Lemma 9 to conclude that  $T \notin \mathcal{F}$ , a contradiction.

**Case 2.**  $t = 8, 9, 10$ .

Since  $t \geq 2^3$ , we may assume that  $(\star)$  holds in  $\mathcal{F}$ . In particular, if  $T \in \mathcal{F}_3$ , then  $|\mathcal{F}(T)| = 7$ .

Let  $\mathcal{D} = \{S \in \binom{[n]}{4} : S \supset T, |\mathcal{F}(S)| = t\}$ . If  $\mathcal{D} = \emptyset$ , then  $\mathcal{F}' = \mathcal{F} - T$  satisfies Property **D** and is thus a  $(4, t)$ -system of size  $|\mathcal{F}| - 1$ , a contradiction. Now suppose that  $|\mathcal{D}| = 1$  and  $\{a\} \cup T$  is the only element of  $\mathcal{D}$ . Since  $t < 11$ , at least one of  $\{a\}, \{a, b\}, \{a, c\}, \{a, d\}$ , say  $\{a\}$ , is not contained in  $\mathcal{F}$ . Let  $\mathcal{F}' = \mathcal{F} - T + \{a\}$ .  $\mathcal{F}'$  satisfies Property **S** trivially. Consider a 4-set  $S \supset T$  of  $[n]$ . If  $S \neq \{a\} \cup T$ , then  $|\mathcal{F}(S)| \geq t + 1$  and  $|\mathcal{F}'(S)| \geq t$ . If  $S = \{a\} \cup T$ , then  $|\mathcal{F}'(S)| = |\mathcal{F}(S)| = t$ . This means that  $\mathcal{F}'$  satisfies Property **D** and consequently  $\mathcal{F}'$  is a  $(4, t)$ -system, a contradiction.

Now we assume that there exist  $a_1, a_2 \in [n]$  such that  $\{a_i\} \cup T \in \mathcal{D}$  for  $i = 1, 2$ . We will show that when  $8 \leq t \leq 10$ , there are two vertices  $v_1, v_2 \in T$  such that  $|\mathcal{F}(\{a_1, a_2, v_1, v_2\})| < t$ , contradicting Property **D**.

Define  $\mathcal{F}_{\{a_i\}}(T) = \mathcal{F}(\{a_i\} \cup T) - \mathcal{F}(T)$  for  $i = 1, 2$ . Since  $|\mathcal{F}(T)| = 7$ , we have  $|\mathcal{F}_{\{a_i\}}(T)| = 1, 2, 3$  for  $t = 8, 9, 10$ , respectively. Using  $(\star)$ , we thus know that  $\{a_i\} \subseteq \mathcal{F}_{\{a_i\}}(T) \subset \mathcal{F}_{\leq 2}$  for every  $t \in \{8, 9, 10\}$ .

- When  $t = 8$ , we have  $\mathcal{F}_{\{a_i\}}(T) = \{\{a_i\}\}$  for  $i = 1, 2$ . Thus  $|\mathcal{F}(\{a_1, a_2, b, c\})| \leq 6 < 8$  for any  $b \neq c \in T$ .
- When  $t = 9$ , we have  $\mathcal{F}_{\{a_1\}}(T) = \{\{a_1\}, \{a_1, c\}\}$  and  $\mathcal{F}_{\{a_2\}}(T) = \{\{a_2\}, \{a_2, d\}\}$ , for not necessarily distinct  $c, d \in T$ . Consequently  $|\mathcal{F}(\{a_1, a_2, b, c\})| \leq 8 < 9$  for some  $b \in T \setminus \{c, d\}$ .
- When  $t = 10$ , we may assume that  $\mathcal{F}_{\{a_1\}}(T) = \{\{a_1\}, \{a_1, b\}, \{a_1, d\}\}$  and  $\mathcal{F}_{\{a_2\}}(T) = \{\{a_2\}, \{a_2, c\}, \{a_2, d\}\}$ , where  $c, b \in T$  are not necessarily distinct. If  $c \neq b$ , then  $|\mathcal{F}(\{a_1, a_2, b, c\})| \leq 8 < 10$ . Otherwise,  $|\mathcal{F}(\{a_1, a_2, b, w\})| \leq 8 < 10$ , where  $w = T \setminus \{c, d\}$ .  $\square$

**Proof of Lemma 7.** Let  $7 \leq t \leq 14$ . We are to show that  $\binom{[n]}{1} \subset \mathcal{F}$ . Suppose instead, say, that  $\{n\} \notin \mathcal{F}$ .

For  $t \geq 8$ , consider a set  $S \in \binom{[n]}{4}$  and  $S \ni n$ . We know that no set from  $\mathcal{F}(S)$  contains  $n$  (otherwise  $(\star)$  forces  $\{n\} \in \mathcal{F}$ ). Thus  $|\mathcal{F}(S)| \leq 7 < t$ , a contradiction to Property **D**.

For  $t = 7$ , consider a set  $T \in \binom{[n-1]}{3}$ . By Property **S** and Property **D**, we have  $|\mathcal{F}(T)| \leq 6$  and  $|\mathcal{F}(\{n\} \cup T)| \geq 7$ . Then there exists a set  $P \in \mathcal{F}(\{n\} \cup T)$  such that  $P \supset n$ . Let  $\mathcal{F}' = \mathcal{F} - P + \{n\}$ . For any  $Y \in \binom{[n]}{3}$  and  $n \in Y$ , we have  $|\mathcal{F}(Y)| \leq 5$  (because  $\{n\}, Y \notin \mathcal{F}$ ). Therefore  $\mathcal{F}'$  satisfies Property **S** and is thus a  $(4, t)$ -system, a contradiction.  $\square$

**Proof of Lemma 8.** We are to show that  $\binom{[n]}{2} \subset \mathcal{F}$  for  $11 \leq t \leq 13$ . Suppose to the contrary, that there exist  $a, b \in [n]$  such that  $\{a, b\} \notin \mathcal{F}$ . Pick any two elements  $v_1, v_2 \in [n] \setminus \{a, b\}$  and consider  $D = \{a, b, v_1, v_2\}$ . Since  $(\star)$  holds, we have  $\{a, b, v_1\}, \{a, b, v_2\} \notin \mathcal{F}$  (otherwise  $\{a, b\} \in \mathcal{F}$ ). Together with  $\{a, b\}$  and  $D$ , this gives us four members of  $(2^D \setminus \emptyset) \setminus \mathcal{F}$ . Consequently  $|\mathcal{F}(D)| \leq 11$ , which contradicts Property **D** when  $t = 12, 13$ . Now assume that  $t = 11$ . Then  $|\mathcal{F}(D)| = 11$  and  $|\mathcal{F}(\{a, v_1, v_2\})| = |\mathcal{F}(\{b, v_1, v_2\})| = 7$ . Let  $\mathcal{F}' = \mathcal{F} - \{a, v_1, v_2\} + \{a, b\}$ .  $\mathcal{F}'$  satisfies Property **S** trivially. To check Property **D**, we consider all the 4-sets  $S$  containing  $\{a, v_1, v_2\}$ . If  $S = \{a, b, v_1, v_2\}$ , then  $|\mathcal{F}'(S)| = |\mathcal{F}(S)| > 11$ . Otherwise,  $S = \{a, v_1, v_2, v_3\}$  for some  $v_3 \in [n] \setminus \{a, b, v_1, v_2\}$ . Since  $|\mathcal{F}(\{a, v_i, v_j\})| = 7$  for any  $i \neq j$ , only  $S$  and  $\{v_1, v_2, v_3\}$  could be absent from  $\mathcal{F}(S)$  and consequently  $|\mathcal{F}(S)| \geq 13$ . We thus have  $|\mathcal{F}'(S)| = |\mathcal{F}(S)| - 1 \geq 13 - 1 > 11$ . Therefore  $\mathcal{F}'$  is a  $(4, 11)$ -system, a contradiction to the optimality of  $\mathcal{F}$ .  $\square$

### 3.2 $m(n, 4, 5)$

In this section we prove that  $m(n, 4, 5) = \binom{n}{2} - 17$ . Before the proof, we introduce the following extensions of the Turán number:

**Definition 10.** A family  $\mathcal{G} \in \binom{[n]}{i}$  is called a *Turán- $i(n, k, t)$ -system* if every  $k$ -set of  $[n]$  contains at least  $t$  members of  $\mathcal{G}$ . The generalized Turán number  $T_i(n, k, t)$  is defined as the minimum size of a *Turán- $i(n, k, t)$ -system*.

Replacing all the instances of  $i$  by  $\geq i$  in the previous paragraph, we obtain the non-uniform Turán number  $T_{\geq i}(n, k, t)$ .

In the proof we will consider  $T_3(k, 4, 1) = \binom{k}{3} - \text{ex}(k, K_4^{(3)})$ . Turán [14] conjectured that

$T_3(k, 4, 1)$  is achieved by the following 3-graph  $\mathcal{H}_k$  (referred to as *Turán's 3-graph*). Partition  $[k]$  into  $A_1 \cup A_2 \cup A_3$ , where  $\lfloor k/3 \rfloor \leq |A_i| \leq \lceil k/3 \rceil$ . The edges of  $\mathcal{H}_k$  are 3-sets which are either contained in some  $A_i$  or contain two vertices of  $A_i$  and one of  $A_{i+1 \pmod{3}}$ . It is known [13] that Turán's conjecture holds for  $k \leq 13$ . For larger  $k$ , the following lower bound of de Caen [1] suffices for our purpose:

$$T_3(k, 4, 1) \geq \frac{k(k-1)(k-3)}{18}. \quad (2)$$

We also need the following simple lemma on  $T_{\geq 1}(n, k, t)$ .

**Lemma 11.** [9]  $T_{\geq 1}(n, k, t) = n - k + t$  for  $1 \leq t \leq k$ .

Let  $\mathcal{F}$  be an optimal  $(4, 5)$ -system with  $A = \{a : \{a\} \in \mathcal{F}\}$ ,  $B = [n] - A$  and assume  $|A| = k$ . By Lemma 5, we may assume that  $\mathcal{F}$  contains no 4-sets. In order to show that  $|\mathcal{F}| \geq \binom{n}{2} - 17$ , our proof consists of three stages described in Section 3.2.1–3.2.3. The proof leads to a construction achieving this bound, which we present in Section 3.2.3 as well.

### 3.2.1 Stage 1

We start with Claim 12 which reflects a rough picture of  $\mathcal{F}$  and in turn implies a (weak) lower bound (4) for  $|\mathcal{F}|$ .

Given two disjoint sets  $C, D \in [n]$ , we write  $\mathcal{F}(C, D) = \{S \in \mathcal{F} : S \cap C \neq \emptyset \text{ and } S \cap D \neq \emptyset\}$ .

**Claim 12.** 1.  $(\mathcal{F}(A))_2$  is a matching in  $A$ .

2.  $\overline{(\mathcal{F}(B))_2}$  contains no matching of size 2 or star with 3 edges.

3.  $|\mathcal{F}(A, B)| \geq (n - k)(k - 2) + |\mathcal{F}_{1,2}(A, B)|$ , where  $\mathcal{F}_{1,2}(A, B) = \{T \in \mathcal{F}_3 : |T \cap A| = 1, |T \cap B| = 2\}$ .

4.  $|(\mathcal{F}(A))_3| \geq k(k - 2)(k - 4)/24$ .

**Proof. Part 1:** Property **S** prevents  $\mathcal{F}(A)$  from containing two adjacent (graph) edges. Thus  $(\mathcal{F}(A))_2$  is a matching.

**Part 2:** We first claim that

$$\text{If } P \in \binom{B}{2} \setminus \mathcal{F} \text{ and } P \subset T, |T| = 3, \text{ then } T \notin \mathcal{F}. \quad (3)$$

In fact, if  $Y$  is 3-set of  $[n]$  such that  $Y \supset P$  and  $|\mathcal{F}(Y)| = 4$ , then  $Y \in \mathcal{F}$ . We may therefore apply Lemma 9 to conclude that  $T \notin \mathcal{F}$ .

If there are  $a, b, c, d \in B$  such that  $\{a, b\}, \{c, d\} \notin \mathcal{F}$ , then  $(\mathcal{F}(\{a, b, c, d\}))_3 = \emptyset$  by (3). Consequently  $|\mathcal{F}(\{a, b, c, d\})| \leq 4$ , a contradiction to Property **D**. Therefore,  $\overline{\mathcal{F}(B)}$  contains no two vertex-disjoint (graph) edges. A similar argument shows that  $\overline{\mathcal{F}(B)}$  contains no star with 3 edges.

**Part 3:** Consider a vertex  $b \in B$  and a 3-subset  $T$  of  $A$ . Since  $\{b\} \notin \mathcal{F}$ ,  $|\mathcal{F}(T)| \leq 4$  and  $|\mathcal{F}(\{b\} \cup T)| \geq 5$ , we have  $|\mathcal{F}(\{b\}, T)| \geq 1$ . Define  $\mathcal{G}_b = \{Y \setminus \{b\} : Y \in \mathcal{F}(\{b\}, A)\}$  for every  $b \in B$ . Then  $\mathcal{G}_b$  is a set system of  $\binom{A}{\leq 2}$  such that every 3-set in  $A$  contains at least one member of  $\mathcal{G}_b$ , in other words,  $\mathcal{G}_b$  is a Turán- $\geq 1(k, 3, 1)$ -system. By Lemma 11, we have  $|\mathcal{H}_b| \geq T_{\geq 1}(k, 3, 1) = k - 2$ . Repeating this for all  $b \in B$ , we have

$$|\{S \in \mathcal{F}(A, B) : |S \cap B| = 1\}| = \sum_{b \in B} |\mathcal{G}_b| \geq (n - k)(k - 2).$$

Consequently  $|\mathcal{F}(A, B)| \geq (n - k)(k - 2) + |\mathcal{F}_{1,2}(A, B)|$ .

**Part 4.** Now we give a crude lower bound for  $(\mathcal{F}(A))_3$ . From Part 1, we know that  $(\mathcal{F}(A))_2$  is a matching  $M = \{\{x_i, y_i\}\}_{i=1}^m$ . Let

$$\mathcal{D} = \{S \in \binom{A}{4} : |S \cap \{x_i, y_i\}| \leq 1 \text{ for every } \{x_i, y_i\} \in M\}.$$

By Property **D**, every 4-set in  $\mathcal{D}$  contains at least one member of  $(\mathcal{F}(A))_3$ . Since  $\mathcal{D}$  is minimal when  $m = \lfloor k/2 \rfloor$ , we may assume that  $m = \lfloor k/2 \rfloor$  when estimating  $(\mathcal{F}(A))_3$  from below. The usual averaging arguments thus give the following lower bound (for even  $k$ , the case when  $k$  is odd yields an even larger bound):

$$(\mathcal{F}(A))_3 \geq \frac{|\mathcal{D}|}{k - 6} = \frac{k(k - 2)(k - 4)(k - 6)}{4!(k - 6)} = \frac{k(k - 2)(k - 4)}{24}. \quad \square$$

The consequence of Claim 12 is the following lower bound.

$$\begin{aligned} |\mathcal{F}| &\geq |(\mathcal{F}(A))_1| + |(\mathcal{F}(A))_3| + |\mathcal{F}(A, B)| + |(\mathcal{F}(B))_2| + |(\mathcal{F}(B))_3| \\ &\geq k + \frac{k(k - 2)(k - 4)}{24} + (n - k)(k - 2) + |\mathcal{F}_{1,2}(A, B)| \\ &\quad + \binom{n - k}{2} - 2 + |(\mathcal{F}(B))_3|. \end{aligned} \tag{4}$$

### 3.2.2 Stage 2

In this stage we first prove Claim 13,  $(\mathcal{F}(A))_2 = \emptyset$ , which not only implies that  $(\mathcal{F}(A))_{\geq 2}$  is a Turán- $_3(k, 4, 1)$ -system, but also makes it possible to find more details about  $\mathcal{F}(A, B)$  and  $\mathcal{F}(B)$ , which are summarized in Claim 14. Claim 13 and 14 together describe the fine structure of  $\mathcal{F}$ . This leads to an improved lower bound (5) for  $|\mathcal{F}|$ .

Let us first sketch the idea behind the proof of Claim 13. Suppose that  $\{a_1, a_2\} \in (\mathcal{F}(A))_2$ . Then at least one of  $B_i = \{b \in B : \{a_i, b\} \notin \mathcal{F}\}$ ,  $i = 1, 2$  has size  $|B|/2$  and consequently either  $|\mathcal{F}_{1,2}(A, B)|$  or  $|(\mathcal{F}(B))_3|$  is at least  $3(n - k)$ . But because of (4),  $|\mathcal{F}|$  is larger than the trivial upper bound  $\binom{n}{2}$ , which is a contradiction.

**Claim 13.**  $(\mathcal{F}(A))_2 = \emptyset$  provided that  $n \geq 160$ .

**Proof.** Note that (4) and  $|\mathcal{F}| \leq \binom{n}{2}$  imply that  $k = O(n^{1/3})$  as  $n \rightarrow \infty$  (in particular, when  $n \geq 20$ ,  $k < n/2$ ).

Suppose instead, that  $\{a_1, a_2\} \in (\mathcal{F}(A))_2$ . Pick a vertex  $b \in B$ . By Property **S**, at most one of  $\{a_1, b\}$  and  $\{a_2, b\}$  is contained in  $\mathcal{F}$ . Without loss of generality, we may assume that  $B$  has a subset  $B_1$  of size  $\frac{n-k}{2}$ , such that  $\{a_1, b\} \notin \mathcal{F}$  for every  $b \in B_1$ . Consider  $\mathcal{T}_{a_1} = \{T \in \mathcal{F}_3 : a_1 \in T, |T \cap B_1| = 2\}$ . If  $|\mathcal{T}_{a_1}| \geq 3(n - k)$ , then (4) implies that (when  $n \geq 30$ ),

$$\begin{aligned} |\mathcal{F}| &\geq k + \frac{k(k-2)(k-4)}{24} + (n-k)(k-2) + \binom{n-k}{2} - 2 + |\mathcal{T}_{a_1}| \\ &\geq \binom{n}{2} + (n-k) + \frac{k(k-2)(k-4)}{24} - \binom{k}{2} + k - 2 \\ &\geq \binom{n}{2} + n - 29 > \binom{n}{2}, \end{aligned}$$

a contradiction to the trivial upper bound that  $|\mathcal{F}| \leq \binom{n}{2}$ , where the third inequality follows from the fact

$$\min_{k \geq 0} \frac{k(k-2)(k-4)}{24} - \binom{k}{2} = -26.125 \quad (\text{achieved by } k = 11).$$

We may therefore assume that  $|\mathcal{T}_{a_1}| < 3(n - k)$ . Let  $\mathcal{P} = \{P \in \binom{B_1}{2} : \{a_1\} \cup P \in \mathcal{T}_{a_1}\}$  and

$\mathcal{T} = \{T \in \binom{B_1}{3} : \binom{T}{2} \cap P = \emptyset\}$ . Then  $|\mathcal{P}| = |\mathcal{T}_{a_1}|$ , and therefore

$$\begin{aligned} |\mathcal{T}| &\geq \binom{(n-k)/2}{3} - |\mathcal{P}| \left( \frac{n-k}{2} - 2 \right) \\ &> \frac{(n-k)(n-k-2)(n-k-4)}{48} - \frac{3}{2}(n-k)^2 \\ &\geq 3(n-k), \quad \text{when } n-k \geq 80 \text{ or } n \geq 160. \end{aligned}$$

On the other hand, we have  $T \in \mathcal{F}$  for every  $T \in \mathcal{T}$  because  $|\mathcal{F}(\{a_1\} \cup T)| \geq 5$  and  $\mathcal{F}(\{a_1\}, T) = \emptyset$ . Consequently  $|(\mathcal{F}(B))_3| \geq |\mathcal{T}| > 3(n-k)$ . Using this lower bound for  $|(\mathcal{F}(B))_3|$  in (4), we obtain  $|\mathcal{F}| \geq k(k-2)(k-4)/24 + \binom{n}{2} + n - 2 > \binom{n}{2}$ , a contradiction.  $\square$

Note that we make no effort to optimize the constant 160 in Claim 13.

With the help of Claim 13, we are able to see the fine structure of  $\mathcal{F}$  as follows.

**Claim 14.** 1.  $(\mathcal{F}(A, B))_3 = \emptyset$ .

2.  $(\mathcal{F}(B))_3 = \emptyset$  and  $|\mathcal{F}(B)| = |(\mathcal{F}(B))_2| \geq \binom{|B|}{2} - 1$ .

3. For every  $a \in A$ , we have  $|\{b \in B : \{a, b\} \in \mathcal{F}\}| \geq n - k - 2$ . Consequently  $|(\mathcal{F}(A, B))_2| \geq k(n - k - 2)$ .

**Proof. Part 1.** Let  $T_0$  be a 3-set of  $[n]$  with  $T_0 \cap A \neq \emptyset$  and  $T \cap B \neq \emptyset$ .

If  $T_0 \cap B = \{b_1, b_2\} \notin \mathcal{F}$ , then  $T_0 \notin \mathcal{F}$  by (3). If  $T_0 \supset \{a, b\} \notin \mathcal{F}$  for some  $a \in A$  and  $b \in B$ , then we consider all the 3-sets  $T \supset P$  such that  $|\mathcal{F}(T)| = 4$ . If  $|T \cap B| = 2$ , then it must be the case that  $T \in \mathcal{F}$ . Otherwise, assume that  $T \cap A = \{a, a'\}$ . Since  $\{a, a'\} \notin \mathcal{F}$  by Claim 13, we also have  $T \in \mathcal{F}$ . We may therefore apply Lemma 9 to conclude that  $T_0 \notin \mathcal{F}$ .

Finally we assume that  $P \in \mathcal{F}$  for every  $P \in \binom{T_0}{2}$  and  $P \not\subset A$ . Either  $|T_0 \cap B| = 2$  or  $|T_0 \cap A| = 2$ , we always have  $|(\mathcal{F}(T_0))_{\leq 2}| = 4$ . Therefore  $T_0 \notin \mathcal{F}$  by Property **S**.

We thus conclude that  $(\mathcal{F}(A, B))_3 = \emptyset$ .

**Part 2.** Suppose instead, that there exists  $T \in (\mathcal{F}(B))_3$ . Then we know that  $\binom{T}{2} \subset \mathcal{F}$  by (3). Let  $\mathcal{D} = \{S \in \binom{[n]}{4} : T \subset S, |\mathcal{F}(S)| = 5\}$ . We may assume that  $\mathcal{D} \neq \emptyset$ , otherwise  $|\mathcal{F}(S)| \geq 6$  for every  $S \supset T$ , and  $T$  could be removed from  $\mathcal{F}$  without hurting Property **D**. Consider a set  $S = \{a\} \cup T \in \mathcal{D}$ . Either  $a \in A$  or  $a \in B$ . If  $a \in A$ , then  $\mathcal{F}(\{a\}, T) = \emptyset$ ; if  $a \in B$ , then  $|\mathcal{F}(\{a\}, T)| = 1$ .

We claim that  $|\mathcal{D}| = 1$ . Suppose instead, that  $\mathcal{D}$  contains two members  $\{a_1\} \cup T$  and  $\{a_2\} \cup T$ . If  $a_1, a_2 \in A$ , then we consider  $S_0 = \{a_1, a_2, b, c\}$  for any two vertices  $b, c \in T$ . From Part 1 we know that  $(\mathcal{F}(S_0))_3 = \emptyset$ . We also have  $\{a_1, a_2\} \notin \mathcal{F}$  by Claim 13. Consequently  $|\mathcal{F}(S_0)| = 3 < 5$ , a contradiction to Property **D**. If  $a_1, a_2 \in B$ , then there are two vertices  $b, c \in T$  such that  $\{a_1, b\}, \{a_2, c\} \notin \mathcal{F}$ . This already contradicts Claim 12 Part 2. Finally, assume that  $a_1 \in A$ ,  $a_2 \in B$  and  $\{a_2, d\} \in \mathcal{F}$  for some  $d \in T$ . Consider  $S_0 = \{a_1, a_2, b, c\}$ , where  $\{b, c\} = T \setminus \{d\}$ . We know that  $(\mathcal{F}(S_0))_3 = \emptyset$  from Part 1 and (3),  $(\mathcal{F}(S_0))_2 \subseteq \{\{a_1, a_2\}, \{b, c\}\}$  from our assumption. Consequently  $|\mathcal{F}(S_0)| = 3 < 5$ , again a contradiction.

Now assume that  $S_0 = \{a\} \cup T$  is the unique element of  $D$ . Let  $\mathcal{F}' = \mathcal{F} - T + \{a, b, c\}$  for any two vertices  $b, c \in T$ .  $\mathcal{F}'$  satisfies Property **S** since  $|\mathcal{F}'(\{a, b, c\})| \leq 3$ .  $\mathcal{F}'$  also satisfies Property **D** because  $|\mathcal{F}'(S)| \geq 6 - 1 = 5$  for every  $S \in \binom{[n]}{4} \setminus S_0$  and  $|\mathcal{F}'(S_0)| = |\mathcal{F}(S_0)| = 5$ .  $\mathcal{F}'$  is thus another optimal  $(4, 5)$ -system. However, if  $a \in A$ , then  $\mathcal{F}'$  contradicts Part 1. If  $a \in B$ , then  $\mathcal{F}'$  contradicts (3), because  $\{a, b, c\} \in \mathcal{F}'$  and at least one of  $\{a, b\}$  and  $\{a, c\}$  is not in  $\mathcal{F}'$ .

We thus conclude that  $(\mathcal{F}(B))_3 = \emptyset$ .

Now it is easy to see why there are no three vertices  $a, b, c \in B$  such that  $\{a, b\}, \{a, c\} \notin \mathcal{F}$ . In such a case, since  $(\mathcal{F}(B))_3 = \emptyset$ , we have  $|\mathcal{F}(\{a, b, c, d\})| \leq 4$  for any  $d \in B \setminus \{a, b, c\}$ , a contradiction to Property **D**. Together with Claim 12 Part 2, we conclude that  $\mathcal{F}(B)$  misses at most one edge on  $B$ .

**Part 3.** Suppose instead, that there exists an  $a \in A$  and  $b_1, b_2, b_3 \in B$  such that  $\{a, b_i\} \notin \mathcal{F}$  for all  $i$ . Part 1 and Part 2 together imply that  $S = \{a, b_1, b_2, b_3\}$  contains no 3-set from  $\mathcal{F}$ . Thus,  $|\mathcal{F}(S)| \leq 4$ , contradicting Property **D**.  $\square$

We now refine (4) by applying Claims 13 and 14:

$$\begin{aligned}
|\mathcal{F}| &= |(\mathcal{F}(A))_1| + |(\mathcal{F}(A))_3| + |(\mathcal{F}(A, B))_2| + |(\mathcal{F}(B))_2| \\
&\geq k + T_3(k, 4, 1) + k(n - k - 2) + \binom{n - k}{2} - 1 \\
&= g(k) + \binom{n}{2} - 1,
\end{aligned} \tag{5}$$

where  $g(k) = T_3(k, 4, 1) - k - \binom{k}{2}$ .

### 3.2.3 Stage 3

In this stage, we complete the proof that  $m(n, 4, 5) = \binom{n}{2} - 17$  by analyzing (5).

Since  $T_3(k, 4, 1)$  is known for  $k \leq 13$ , we compute  $g(k)$  exactly for  $0 \leq k \leq 11$  and obtain that  $\min_{0 \leq k \leq 11} g(k) = g(7) = g(8) = -16$ . For  $k \geq 12$ , using the inequality (2), we have  $g(k) \geq k(k-1)(k-3)/18 - k - \binom{k}{2} \geq -12$ . Putting these together, we have

$$\min_{k \geq 0} g(k) = g(7) = g(8) = -16 \quad (6)$$

Applying (6) to (5), we obtain that  $|\mathcal{F}| \geq \binom{n}{2} - 17$ .

Claims 13, 14 and (6) lead us to the following construction, which gives a  $(4, 5)$ -system of cardinality  $\binom{n}{2} - 17$ .

**Construction 1:** Partition  $[n]$  into  $A \cup B$ , where  $|A| = k = 7$  or  $8$ . Let  $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3$ , where  $\mathcal{F}_1$ ,  $\mathcal{F}_2$  and  $\mathcal{F}_3$  are defined as follows:

- $\mathcal{F}_1 = \{\{a\} : a \in A\}$ .
- Let  $M$  be a union of  $k$  disjoint copies of  $P_2$  (paths of length 2) whose middle vertices are in  $A$  and end vertices make up a subset  $B_0$  of  $B$ . Let  $\mathcal{F}_2 = \binom{B}{2} \setminus \{e\} \cup (A \times B) \setminus M$  for some  $e \in B \setminus B_0$ .
- $\mathcal{F}_3$  is the edge set of the Turán 3-graph  $\mathcal{H}_k$  on  $A$ .

We thus conclude that  $m(n, 4, 5) = \binom{n}{2} - 17$ . □

### 3.3 $m(n, 4, 6)$

Let  $\mathcal{F}$  be an optimal  $(4, 6)$ -system. Define  $A$ ,  $B$ ,  $\mathcal{F}(A)$ ,  $\mathcal{F}(B)$ ,  $\mathcal{F}(A, B)$  as in Section 3.2 and assume that  $|A| = k$ . We first define another threshold function.

**Definition 15.** A  $(\widehat{4, 2})$ -system of  $[n]$  is a set system  $\mathcal{G} \subseteq \binom{[n]}{2} \cup \binom{[n]}{3}$  such that every 4-set of  $[n]$  contains at least two members of  $\mathcal{G}$  and every 3-set of  $[n]$  contains at most two members of  $\mathcal{G}$ . Let  $\hat{m}(n, 4, 2)$  denote the minimum size of such a set system.

The lower bound  $|\mathcal{F}| \geq \binom{n}{2} - 190$  in Theorem 3 is the consequence of the following claim, whose proof is postponed to the end of this section.

**Claim 16.**  $\mathcal{F}$  has the following properties.

1. There exists no  $T \in \mathcal{F}_3$  which contains  $a \in A$  and  $b \in B$  such that  $\{a, b\} \notin \mathcal{F}$ .
2.  $\mathcal{F}(B) = \binom{B}{2}$ .
3.  $|\mathcal{F}(A, B)| \geq k(n - k) - 4$ .
4.  $|\mathcal{F}_{\geq 2}(A)| \geq \hat{m}(k, 4, 2) \geq 2\binom{k}{7}/\binom{k-3}{4} = k(k-1)(k-2)/105$ .

Claim 16 also suggests a general way to construct  $(4, 6)$ -systems of  $[n]$ : Partition  $[n]$  into  $A \cup B$  with  $|A| = k$  for any  $0 \leq k \leq n$ . Let  $\mathcal{F} = \mathcal{F}(A) \cup \mathcal{F}(A, B) \cup \mathcal{F}(B)$ , with  $\mathcal{F}(B) = \binom{B}{2}$ ,  $\mathcal{F}(A, B) = A \times B$  and  $\mathcal{F}(A) = (\mathcal{F}(A))_1 \cup (\mathcal{F}(A))_{\geq 2}$ , where  $(\mathcal{F}(A))_1 = \{\{a\} : a \in A\}$  and  $(\mathcal{F}(A))_{\geq 2}$  is a  $(4, 2)$ -system on  $A$ .

In particular, the following construction gives a  $(4, 6)$ -systems of  $[n]$  of size  $\binom{n}{2} - 5$ .

**Construction 2:** Let  $k = 8$  and  $A, B, \mathcal{F}(B), \mathcal{F}(A, B)$  and  $(\mathcal{F}(A))_1$  are defined as above. We construct  $(\mathcal{F}(A))_{\geq 2}$  as follows.

- Suppose that  $A = A_1 + A_2$ , where  $A_1 = \{u_1, u_2, u_3, u_4\}$  and  $A_2 = \{v_1, v_2, v_3, v_4\}$ . Let  $E_0 = \{\{u_1, v_1\}, \{u_1, v_2\}, \{u_2, v_3\}, \{u_3, v_3\}, \{u_4, v_4\}\}$ . Let  $(\mathcal{F}(A))_2 = (A_1 \times A_2) - E_0$ .
- $(\mathcal{F}(A))_3 = \{\{u_1, u_2, u_3\}, \{u_2, u_3, u_4\}, \{v_1, v_2, v_3\}, \{v_1, v_2, v_4\}\}$ .

**Proof of Theorem 3 for  $t = 6$ .** The upper bound  $m(n, 4, 6) \leq \binom{n}{2} - 5$  follows from Construction 2. Claim 16 gives the lower bound for  $|\mathcal{F}| = m(n, 4, 6)$  as follows.

$$\begin{aligned}
|\mathcal{F}| &= |(\mathcal{F}(A))_1| + |(\mathcal{F}(A))_{\geq 2}| + |\mathcal{F}(A, B)| + |\mathcal{F}(B)| \\
&\geq k + \frac{k(k-1)(k-2)}{105} + k(n-k) - 4 + \binom{n-k}{2} \\
&\geq \binom{n}{2} - 190,
\end{aligned} \tag{7}$$

where the last inequality follows from  $\min_{k \geq 0} k + k(k-1)(k-2)/105 - \binom{k}{2} = -186$  (achieved by  $k = 35$ ).  $\square$

**Remark:** Actually, we almost determine  $m(n, 4, 6)$  exactly in terms of  $\hat{m}(k, 4, 2)$  (off only by 4). To see this, replace  $k(k-1)(k-2)/105$  by  $\hat{m}(k, 4, 2)$  in (7) and get an upper bound

following the general construction:

$$k + \hat{m}(k, 4, 2) + k(n - k) + \binom{n - k}{2} - 4 \leq |\mathcal{F}| \leq k + \hat{m}(k, 4, 2) + k(n - k) + \binom{n - k}{2},$$

for some  $k \geq 0$ . The knowledge of  $\hat{m}(k, 4, 2)$  for small values of  $k$  may lead to the final settlement of  $m(n, 4, 6)$ .

**Proof of Claim 16.**

**Part 1:** Suppose that there are two vertices  $a \in A$  and  $b \in B$  such that  $\{a, b\} \notin \mathcal{F}$ . For a 3-set  $T \supset \{a, b\}$ , if  $|\mathcal{F}(T)| = 5$ , then  $T \in \mathcal{F}$ . We may therefore apply Lemma 9 to conclude that  $T \notin \mathcal{F}$  for every 3-set  $T \supset \{a, b\}$ .

**Part 2:** We first claim that (3) holds in  $\mathcal{F}$ . In fact, when  $P \in \binom{B}{2}$ , we have  $\{T \in \binom{[n]}{3} : P \subset T, |\mathcal{F}(T)| = 5\} = \emptyset$ . Then we can apply Lemma 9 to obtain (3).

Next, we show that if there exists a set  $T \in (F(B))_3$ , then we obtain a contradiction. The proof is similar to that of Claim 14 Part 2. First, we claim that  $\mathcal{D} = \{S \in \binom{[n]}{4} : T \subset S, |\mathcal{F}(S)| = 6\}$  has exactly one member. Suppose instead, for example, that  $\mathcal{D}$  contains  $S_1 = \{a\} \cup T$  and  $S_2 = \{b\} \cup T$  for some  $a \in A$  and  $b \in B$  (other cases are similar). It means that there are exactly two sets from  $\mathcal{F}(S_1)$ ,  $\mathcal{F}(S_2)$  which contain  $a$ ,  $b$ , respectively. From Part 1, we know that these two sets in  $S_1$  must be  $\{a\}$  and  $\{a, b_1\}$  and the two sets in  $S_2$  must be  $\{b, b_2\}$  and  $\{b, b_3\}$ , where  $b_1, b_2, b_3 \in T$ . Assume that, for example,  $b_1, b_2, b_3$  are all distinct. Consider  $S_3 = \{a, b, b_1, b_2\}$ . It is easy to see that  $|\mathcal{F}(S_3)| \leq 5$ , a contradiction to Property **D**. Now assume that  $\{a\} \cup T$  is the unique member of  $\mathcal{D}$ , for example, for some  $a \in A$ . Let  $\mathcal{F}' = \mathcal{F} - T + \{a, b_1, b_2\}$ . It is easy to see that  $\mathcal{F}'$  is an optimal  $(4, 6)$ -system. However, since  $\{a, b_1, b_2\} \in \mathcal{F}'$  and  $\{a, b_2\} \notin \mathcal{F}'$ , this contradicts Part 1.

Since  $(F(B))_3 = \emptyset$ ,  $\binom{B}{2} \subset \mathcal{F}$  follows from Property **D**. Consequently  $\mathcal{F}(B) = \binom{B}{2}$ .

**Part 3:** We first show that for every  $a \in A$ , there is at most one vertex  $b \in B$  such that  $\{a, b\} \notin \mathcal{F}$ . Suppose instead, that  $\{a, b_1\}, \{a, b_2\} \notin \mathcal{F}$  for some  $a \in A$ . Consider  $S = \{a, b_1, b_2, c\}$  for any vertex  $c \in B \setminus \{b_1, b_2\}$ . From Part 1 and 2 we know that  $(\mathcal{F}(S))_3 = \emptyset$ . Consequently  $|\mathcal{F}(S)| \leq 5$ , a contradiction to Property **D**. Second, for each  $b \in B$ , there are at most two vertices  $a_1, a_2 \in A$  such that  $\{a_i, b\} \notin \mathcal{F}$  for  $i = 1, 2$ . Suppose instead, that there are three vertices  $a_1, a_2, a_3 \in A$  such that  $\{a_i, b\} \notin \mathcal{F}$ . Since  $\mathcal{F}(\{a_1, a_2, a_3\}, \{b\}) = \emptyset$ , we have  $\mathcal{F}(\{a_1, a_2, a_3\}) = \mathcal{F}(\{b, a_1, a_2, a_3\})$ , which either violates Property **D** or Property **S**.

Now consider the bipartite graph  $G$  whose edge set is  $(A \times B) - (\mathcal{F}(A, B))_2$ . By the argument

in the previous paragraph,  $G$  consists of vertex-disjoint edges or 2-paths whose centers are in  $B$ . On the other hand, two independent edges  $\{a_i, b_i\}$ ,  $i = 1, 2$  in  $G$  (where  $a_i \in A$ ) imply that  $\{a_1, a_2\} \in \mathcal{F}$  (by Property **D**). If  $G$  contain 3 independent edges  $\{a_i, b_i\}$ , then  $|\mathcal{F}(\{a_1, a_2, a_3\})| = 6$ , a contradiction to Property **S**. Therefore  $G$  has at most 4 edges which are from two disjoint 2-paths.

**Part 4.** Clearly  $\mathcal{F}_{\geq 2}(A)$  is a  $(\widehat{4, 2})$ -system of  $[k]$ . Let us count the number of triples in a  $(\widehat{4, 2})$ -system  $\mathcal{G}$  of  $[k]$ . The following lemma implies that every 7-set of  $[k]$  contains at least two triples from  $\mathcal{G}$ . We omit its proof because it is an easy case analysis.

**Lemma 17.** *Every  $(\widehat{4, 2})$ -system of  $[7]$  contains at least two triples.* □

Using Lemma 17 and the averaging argument, we have  $\hat{m}(k, 4, 2) \geq 2 \binom{k}{7} / \binom{k-3}{4}$ . □

## 4 Acknowledgements

The authors thank Gy. Turán for introducing them to the function  $m(n, k, t)$  and helpful discussions.

## References

- [1] D. de Caen, Extension of a theorem of Moon and Moser on complete subgraphs. *Ars Combin.* **16** (1983), 5–10.
- [2] F. Chung, L. Lu, An upper bound for the Turán number  $t_3(n, 4)$ . *J. Combin. Theory Ser. A* **87** (1999), no. 2, 381–389.
- [3] P. Erdős, A. Rényi, On a problem in the theory of graphs. (Hungarian) *Magyar Tud. Akad. Mat. Kutat Int. Közl.* **7** 1962 623–641 (1963).
- [4] P. Erdős, M. Simonovits, Compactness results in extremal graph theory. *Combinatorica* **2** (1982), no. 3, 275–288.
- [5] P. Frankl, Z. Füredi, An exact result for 3-graphs, *Discr. Math.* **50** (1984), 323–328.
- [6] Z. Füredi, R. H. Sloan, K. Takata, Gy. Turán, On set systems with a threshold property, submitted.

- [7] Kővari, T.; Sós, V. T.; Turán, P. On a problem of K. Zarankiewicz. *Colloquium Math.* **3**, (1954). 50–57
- [8] D. Mubayi, On hypergraphs with every four points spanning at most two triples, submitted
- [9] D. Mubayi, Y. Zhao, Non-Uniform Turán-Type problems, submitted
- [10] S. Jukna, Computing threshold functions by depth-3 threshold circuits with smaller thresholds of their gates, *Inf. Proc. Lett.* **56** (1995), 147–150
- [11] R. H. Sloan, K. Takata, Gy. Turán, On frequent sets of Boolean matrices, *Annals of Mathematics and Artificial Intelligence* **24** (1998) 193–209
- [12] J. H. Spencer, Maximal consistent families of triples, *J. Combin. Theory* **5** (1968), 1-8.
- [13] T. H. Spencer, On the size of independent sets in hypergraphs. Coding theory, design theory, group theory (Burlington, VT, 1990), 263–273, Wiley-Intersci. Publ., Wiley, New York, 1993.
- [14] P. Turán, On an extremal problem in graph theory. *Mat. Fiz. Lapok* **48** (1941), pp. 436-452.